

- (c) Providing large number of options for learner to choose major and minor subjects in core and allied areas as well as skill and electives in areas of interest.
- (d) Make research an integral part of learning through Projects, Assignments, Visits; guest/expert talks value added courses, extracurricular/Co-Curricular activities with credits added for these.
- (e) Character building and social connect as part of the curriculum in words and spirit.
- (f) Collaboration with industries and other institutes in India and abroad to keep the learner abreast with outside world.

## REFERENCES

- [1] Mc Cowan, T (2019) Higher Education for and beyond the sustainable goals. Springer Nature.
- [2] National Education Policy 2020. Ministry of Human Resource Development Govt of India.
- [3] Fostering Social Responsibility in HEIs. University Grant Commission. [https://www.ugc.ac.in/e-book/UNNAT % 20 HBARAT %ABHIYAN](https://www.ugc.ac.in/e-book/UNNAT%20HBARAT%ABHIYAN). Pdf.
- [4] Ministry of Education, India website <https://www.education.govt.in/research>

## Wastewater Treatment by Low Cost Technology

Amitabh Sharma

Dept. of Chemistry, Dr. C.V. Raman University, Bilaspur (C.G.) India.

### ABSTRACT

*Nowadays many water resources are polluted by anthropogenic sources including household and agricultural waste and industrial processes. Public concern over the environmental impact of wastewater pollution has increased. Several conventional wastewater treatment techniques, i.e. chemical coagulation, adsorption, activated sludge, have been applied to remove the pollution, however there are still some limitations, especially that of high operation costs. The use of aerobic waste water treatment as a reductive medium is receiving increased interest due to its low operation and maintenance costs. In addition, it is easy-to-obtain, with good effectiveness and ability for degrading contaminants. This paper reviews the use of waste water treatment technologies to remove contaminants from wastewater such as halogenated hydrocarbon compounds, heavy metals, dyes, pesticides, and herbicides, which represent the main pollutants in wastewater.*

**Key Words:** Sewage Treatment, Technologies and domestic wastewater.

### I INTRODUCTION

A supply of clean water is an essential requirement for the establishment and maintenance of diverse human activities. Water resources provide valuable food through aquatic life and irrigation for agriculture production. However, liquid and solid wastes produced by human settlements and industrial activities pollute most of the water sources throughout the world. Due to massive worldwide increases in the human population, water will become one of the scarcest resources in the 21st century (Day D., 1996). In the year 2015 the majority of the global population (over 5 billion) will live in urban environments (UN, 1997). By the year 2015, there will be 23 megacities with a population of over 10 million each, 18 of which will exist in the developing world (Black, 1994). Central to the urbanization phenomena are the problems associated with providing municipal services and water sector infrastructure, including the provision of both fresh water resources and sanitation services. Currently, providing housing, health care, social services, and access to basic human needs infrastructure, such as clean water and the disposal of effluent, presents major challenges to engineers, planners and politicians (Black, 1994; Giles and Brown, 1997).

As human numbers increase, greater strains will be placed on available resources and pose even greater threat to environmental sources. A report by the Secretary-General of the United Nations Commission on Sustainable Development (UNCSD, 1997) concluded that there is no sustainability in the current uses of fresh water by either developing or developed nations, and that worldwide, water usage has been growing at more than three times the world's population increase, consequently leading to widespread public health problems, limiting economic and agricultural development and adversely affecting a wide range of ecosystems. Although India occupies only 3.29 million km<sup>2</sup> geographical area, which forms 2.4% of the world's

land area, it supports over 15% of world's population. The population of India as of March 1, 2001 was 1,027,015,247 persons (Census, 2001).

India also has a livestock population of 500 million, which is about 20% of world's total livestock. However, total annual utilizable water resources of the country are 1086 km<sup>3</sup> which is only 4% of world's water resources (Kumar et al., 2005). Total annual utilizable resources of surface water and ground water are 690 and 396 km<sup>3</sup>, respectively (Ministry of Water Resources, 1999). Consequent to rapid growth in population and increasing water demand, stress on water resources in India is increasing and per capita water availability is reducing day by day. In India per capita surface water availability in the years 1991 and 2001 were 2300 m<sup>3</sup> (6.3 m<sup>3</sup>/day) and 1980 m<sup>3</sup> (5.7 m<sup>3</sup>/day) respectively and these are projected to reduce to 1401 and 1191 m<sup>3</sup> by the years 2025 and 2050, respectively (Kumar et al., 2005). Total water requirement of the country in 2050 is estimated to be 1450 km<sup>3</sup> which is higher than the current availability of 1086 km<sup>3</sup>. Much of the wastes of civilization enter water bodies through the discharge of waterborne waste from domestic, industrial and non-point sources carrying unwanted and unrecovered substances (Welch, 1992).

Although the collection of wastewater dates back to ancient times, its treatment is a relatively recent development dating from the late 1800s and early 1900s (Chow et al., 1972). Modern knowledge of the need for sanitation and treatment of polluted waters however, started with the frequently cited case of John Snow in 1855, in which he proved that a cholera outbreak in London was due to sewage contaminated water obtained from the Thames River (Cooper, 2001). In developed nations, treatment and discharge systems can sharply differ between countries and between rural and urban users, with respect to urban high income and urban low-income users (Doorn et al., 2006). The most common wastewater treatment methods in developed countries are centralized aerobic wastewater treatment plants and lagoons for both domestic and industrial wastewater.

The degrees of wastewater treatment vary in most developing countries. Domestic wastewater may be treated in centralized plants, pit latrines, septic systems or disposed of in unmanaged lagoons or waterways, via open or closed sewers (UNEP, 2002). In some cases industrial wastewater is discharged directly into water bodies, while major industrial facilities may have comprehensive inplant treatment (Carter et al., 1999; Doorn et al., 2006). Modern civilization, armed with rapidly advancing technology and fast growing economic system is under increasing threat from its own activities causing water pollution, (Singh et al. (1989). India is the seventh largest country in the world with a total landmass of 3.29 million sq. km, population over 1 billion, 29% of which live in urban areas spread over 5162 towns. With enormous natural resources and growing economy India is the second largest pool of technical and scientific personnel in the world. Pollution from small size industries (SSIs) puts the Indian regulators in front of a difficult arbitrage between economic development and environmental sustainability.

The uncontrolled growth in urban areas has made planning and expansion of water and sewage systems very difficult and expensive (Looker, 1998).

Aerobic activated sludge reactors have been used on a limited scale as bio-scrubbers for the treatment of odorous air (Bowker, 2000). Despite numerous positive reports from full scale applications in North America, little data are available on the actual performance of these systems with wide ranging concerns on reduction of settling efficiency due to changes in filamentous organisms and bacterial flocks (Burgess et al. 2001). These concerns are alleviated in MBRs where gravitational settling of the microbial solution is replaced by physical filtration. Also, the diffusion and bioconversion of odorous gases are a function of contact time, bubble size, and reactor configuration (Burgess et al. 2001). Submerged MBRs incorporate the membrane unit within the bioreactor and rely on gas and liquid scouring to clean the membrane surface. Since modern livestock operations are equipped with blowers and ventilation systems, booster fans could be added to increase outflow pressure. This concept was explored in past research efforts when biofilter beds (compost and wood chips) were tested for odour removal (Mann et al. 2002).

## II STATUS OF WASTEWATER IN INDIA

The total wastewater generated by 299 class-1 cities is 16,652.5 MLD. Out of this, about 59% is generated by 23 metro cities. The state of Maharashtra alone contributes about 23%, while the Ganga river basin contributes about 31% of the total wastewater generated in class-1 cities. Only 72% of the total treated wastewater generated is collected. Out of 299 class-1 cities, 160 cities have sewerage system for more than 75 percent of population and 92 cities have more than 50 percent of population coverage. On the whole 70% of total population of class-1 cities is provided with sewerage facility, compared to 48% in 1988. The type of sewerage system is either open or closed or piped. The main objective of this study was to perform a review of the treatment of domestic sewage using the aerobic sludge to ensure effective discharge and/or re-use/recycling.

## III WASTEWATER TREATMENT IN INDIA

Out of 16,662.5 MLD of wastewater generated, only 4037.2 mld (24 %) is treated before release, the rest (i.e. 12,626.30 MLD) is disposed of untreated. Twenty-seven cities have only primary treatment facilities and only forty-nine have primary and secondary treatment facilities.

- (a) **Need of sewage treatment:** Wastewater treatment involves breakdown of complex organic compounds in the wastewater into simpler compounds that are stable and nuisance-free, either physico-chemically and/or by using micro-organisms (biological treatment). The adverse environmental impact of allowing untreated wastewater to be discharged in groundwater or surface water bodies and or lands are as follows:
- (i) The decomposition of the organic materials contained in wastewater can lead to the production of large quantities of malodorous gases.
  - (ii) Untreated wastewater (sewage) containing a large amount of organic matter, if discharged into a river / stream, will consume the dissolved oxygen for satisfying the Biochemical Oxygen Demand (BOD) of wastewater and thus deplete the dissolved oxygen of the stream, thereby causing fish kills and other undesirable effects.
  - (iii) Wastewater may also contain nutrients, which can stimulate the growth of aquatic plants and algal blooms, thus leading to eutrophication of the lakes and streams.
  - (iv) Untreated wastewater usually contains numerous pathogenic, or disease causing microorganisms and toxic compounds, that

dwelling in the human intestinal tract or may be present in certain industrial waste. These may contaminate the land or the water body, where such sewage is disposed.

For the above-mentioned reasons the treatment and disposal of wastewater, is not only desirable but also necessary.

#### IV INDUSTRIAL, MUNICIPAL AND DOMESTIC REUSE OF WASTEWATER

Municipal uses of treated wastewater include the irrigation of road plantings, parks, playgrounds, golf courses and toilet flushing etc. (Bouwer, 1993). Industrial reuses of wastewater include cooling systems, agricultural uses (irrigation and aquaculture), the food processing industry and other highrate water uses (Bouwer, 1993b; Khouri *et al.* 1994; Asano and Levine, 1996). In Middle Eastern countries, where water is scarce, dual distribution systems will, in the near future, provide high quality, treated effluents for toilet flushing to hotels, office buildings, etc. (Shelef and Azov, 1996). In India, wastewater is currently being used for irrigation, gardening, flushing, cooling of air conditioning systems, as a feed for boilers, and as process water for industries (Chawathe and Kantawala, 1987). In China, national policy has been developed that promotes the development of water-efficient technologies, and encourages the reuse of reclaimed municipal wastewater in agriculture first, and then for industrial and municipal uses (Zhongxiang and Yi, 1991).

#### V CONCLUSION

This report is a review of variety of options that may be employed in the treatment, recovery and reuse of wastewater. It is apparent that a variety of options are feasible for use in the developing world and even more apparent that many low-technology options can be mixed and matched for very high efficiencies. Natural treatment technologies are attracting a significant level of interest by environmental managers. Natural treatment technologies are considered viable because of their low capital costs, their ease of maintenance, their potentially longer life-cycles and their ability to recover a variety of resources including: treated effluent for irrigation, organic humus for soil amendment and energy in the form of biogas. This report examined emergent issues and technological options related to the scale of collection and treatment systems. There is increasing

Momentum developing behind the notion that recycling loops, from point of generation (*e.g.*, the household) to point of treatment and reuse must be shortened.

#### REFERENCES

- [1] Asano, T. and A. D. Levine (1996). Wastewater reclamation, recycling, and reuse: Past, present and future. *Water Science and Technology*, 33: 1-14.
- [2] Boller, M. (1997). Small wastewater treatment plants - A challenge to wastewater engineers. *Water Science and Technology*, 35: 1-12.
- [3] Bouwer, H. (1993a). From sewage to zero discharge. *European Water Pollution Control*, 3: 9-16.
- [4] Burgess, J.E., S.A. Parsons and R.M. Stuetz (2001). Developments in odour control and waste gas treatment biotechnology: A review. *Biotechnology Advances*, 19: 35-63.
- [5] Carter, C.R., S.F. Tyrrel and P. Howsam (1999). Impact and sustainability of community water supply and sanitation programmes in developing countries. *Journal of the Chartered Institution of Water and Environmental Management*, 13: 292-296.
- [6] Census of India (2001): Analysis and Articles on Population and Literacy Rates, Office of the Registrar General, India, Ministry of Home Affairs.
- [7] Chawathe, S. D. and D. Kantawala (1987). Reuse of water in city planning. *Water Supply*, 15: 17-23.
- [8] Cooper, P.F. (2001). Historical aspects of wastewater treatment. In *Decentralized sanitation and reuse concepts, systems and Implementation*. Eds., Lens P., Zeeman G., and G. Lettinga. IWA Publishing. London. pp. 11-38.
- [9] Deb, P.K., A.J. Rubin, A.W. Launder and K.H. Mancy (1966). Removal of COD from wastewater by fly ash. In *proceeding of the 1966. 21st Indiana Waste conference*. D.E. Bloodygood Ed. Published by Purdue University, W. Lafayette, Indiana. pp: 848 – 860.
- [10] Doorn, M.R.J., S. Towprayoon, S. Maria, M. Vieira, W. Irving, C. Palmer, R. Pipatti and C.Wang, (2006). Wastewater treatment and discharge. In *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. WMO, UNEP. pp. 5: 1-6

- [11] Frijns, J. and M. Jansen (1996). Institutional requirements for appropriate wastewater treatment systems. In A. Balkema, H. Aalbers and E. Heijndermans (Eds.), Workshop on sustainable municipal waste water treatment systems, Leusdan, the Netherlands. ETC in cooperation with WASTE. 12-14 November, 1996. pp: 54-66.
- [12] Harada, H., M. Tandulkar and A. Ohashi, (2007). Performance comparison of a pilot-scale UASB and DHS system and activated sludge process for the treatment of municipal wastewater. *Water Research*, 132: 166-172.
- [13] Harada, H., M. Tandulkar, S. Uemura and A. Ohashi (2005). A low-cost municipal sewage treatment system with a combination of UASB and the Fourth Generation Down flow Hanging Sponge (DHS) reactors. *Water Science and Technology*, 52: 323-329.
- [14] Khouri, N., J.M. Kalbermatten and C. Bartone (1994). The reuse of wastewater in agriculture: A guide for planners. UNDP - World Bank Water and Sanitation Program. Washington, DC: The World Bank.
- [15] Looker, N. (1998). *Municipal Wastewater Management in Latin America and the Caribbean*, R.J. Burnside International Limited, Published for Roundtable on Municipal Water for the Canadian Environment Industry Association. *International Journal of Engineering Research & Technology (IJERT)* Vol. 1 Issue 5, July – 2012
- [16] Nelson, M. and C.F. Guarino (1969). The use of fly ash in municipal waste treatment. *Journal of the Water Pollution Control Federation*, 41: 1905-1911.
- [17] Tajrishi, M. and A. Abrishamchi (2005). *Integrated Approach to Water and Wastewater Management for Tehran, Iran*, "Water Conservation, Reuse and Recycling: Proceedings of an Iranian-American Workshop 2005", the National Academies Press, Washington, D.C.
- [18] Yu, H., Tay, J. and F. Wilson (1997). A sustainable municipal wastewater treatment process for tropical and subtropical regions in developing countries. *Water Science and Technology*, 35: 191-198.